Capitainer B, HbA1c

Manufacturer Capitainer AB

Supplier in Denmark You Do Bio

Supplier in Sweden Same as the manufacturer

Supplier in Norway Same as the manufacturer

Launched in Scandinavia February 2020



Aim

To assess the analytical performance and user-friendliness of Heamoglobin A1c (HbA1c) measurements of capillary blood, collected with Capitainer B sampling card, analysed on Roche cobas pro with Tina-quant. Sample collection was performed by the intended users (health care personnel and lay persons), and measurements of HbA1c were performed by biomedical laboratory scientists. The first evaluation showed poor accuracy, therefore, a second evaluation was carried out after the manufacturer introduced changes to the pre-analytical procedure, as well as shortening the recommended stability time of the Capitainer B sampling card, from 14 to 5 days after sampling. The second evaluation included only Capitainer B samples collected by lay persons with diabetes, recruited at primary health care centers (PHCCs).

Performance specifications	Results	Conclusions
Repeatability	First evaluation	Fulfilled when samples were
CV ≤3,0 %	CV 1,4 – 2,8 %	collected by health care personnel.
	Second evaluation	Fulfilled when samples were
	CV 0,8 – 1,3 %	collected by lay persons.
Accuracy	First evaluation	Not fulfilled when samples were
\geq 95 % of individual results within \pm 3,0 mmol/mol at HbA1c concentration $<$ 35,3 mmol/mol and \pm 8,5 %	74,0 %	collected by health care personnel.
at HbA1c concentration ≥35,3 mmol/mol compared	Second evaluation	Not fulfilled when samples were
to the average result of the comparison method.	89,0 %	collected by lay persons.
User-friendliness	First and second evaluation	Fulfilled
A total rating of "Satisfactory"	The user-friendliness was rated	
	satisfactory.	

Additional information

Auditional information		
Participants and evaluation sites First evaluation	113 persons with diabetes recruited at a hospital laboratory and 92 persons with diabetes recruited at three primary health care centres.	
Second evaluation	115 persons with diabetes recruited at seven primary health care centres.	
Evaluated method Both evaluations	HbA1c on capillary blood samples collected with Capitainer B sampling card, analysed on cobas pro c503 with Tina-quant.	
Comparison method Both evaluations	cobas pro c503 with Tina-quant. HbA1c on venous whole blood samples.	
Bias Both evaluations	When samples were collected by health care personnel in the first evaluation, a positive bias of 2,1 to 3,0 mmol/mol was observed between the evaluated method and the comparison method. When samples were collected by lay persons in the second evaluation, a positive bias of 0,7 to 2,1 mmol/mol was observed between the methods.	

A letter with comments from Capitainer AB is attached to the report.

Further information about the evaluation and the organisation of SKUP can be found on www.skup.org.

This summary is also published in Danish, Norwegian and Swedish at www.skup.org.

Copyright © 2025 SKUP. SKUP is a co-operative commitment of DEKS in Denmark, Equalis in Sweden and Noklus in Norway. The secretariat is located at Noklus in Bergen, Norway.